


Only use Azelastin-COMOD if the sealing of the folding carton was intact before the first use.

Wash your hands.


Take a clean tissue and gently wipe around your eyes to remove any moisture.



**Step 1:**


Remove the cap before use.

Before the first application please turn the Azelastin-COMOD-bottle with its tip down and press onto the bottom of the bottle until the first drop appears at the nozzle. Now the bottle is ready for further use.



**Step 2:**

Hold the bottle upside down with the thumb on the shoulder of the bottle and the other fingers on the bottom of the bottle.



**Step 3:**

Support the hand holding the COMOD<sup>®</sup>-bottle as shown with the other hand.




**Step 4:**

Lean your head back slightly, gently pull down your lower eyelid and apply a strong and quick pressure on the middle bottom of the bottle. This activates the mechanism for the release of one drop. Due to the special COMOD<sup>®</sup> valve system, size and speed of each drop are the same no matter how much pressure is applied. Slowly close your eyes to allow the fluid to spread evenly on the eye surface.



**Step 5:**

After instilling Azelastin-COMOD release the lower eye lid and press with your finger on the corner of the eye closest to the nose (see picture). While doing this blink several times with your eyes to distribute the eye drops on the eye ball. Remove surplus eye drops afterwards.



**Step 6:**

Replace cap immediately after use. Take care that the dropper tip is dry.

**Duration of treatment**

If possible, you should use Azelastin-COMOD regularly until your symptoms have disappeared.

Do not use Azelastin-COMOD for more than 6 weeks.